

**S10 Table. Pediatric Population Characteristics**

	Entire Cohort	NNC <sup>1</sup>	CNS <sup>1</sup>	PNS <sup>1</sup>	P-value <sup>2</sup>
<b>All Patients, N (% of cohort)</b>	2198 (100%)	2019 (91.9%)	163 (7.4%)	16 (0.73%)	
<b>Sex, N (%)</b>					.074
Female	1026	954 (47.3%)	62 (38%)	10 (62.5%)	
Male	1172	1065 (52.7%)	101 (62%)	6 (37.5%)	
<b>Age, N (%)</b>					
0-2	769	747 (37.3%)	22 (15.8%)	0 (0%)	<.001*
3-5	275	251 (12.5%)	24 (17.3%)	0 (0%)	.283
6-11	403	368 (18.4%)	33 (23.7%)	2 (16.7%)	.763
12-17	707	637 (31.8%)	60 (43.2%)	10 (83.3%)	.039*
<b>Race, N (%)</b>					
American Indian	1	1 (0%)	0 (0%)	0 (0%)	.957
Asian	61	52 (2.6%)	9 (5.8%)	0 (0%)	.119
Black	184	166 (8.3%)	16 (10.4%)	2 (12.5%)	.763
Hawaiian/Pacific Islander	1	1 (0%)	0 (0%)	0 (0%)	.957
Hispanic/Latino	20	20 (1%)	0 (0%)	0 (0%)	.502
White	665	608 (30.2%)	47 (30.5%)	10 (62.5%)	.048*
Other / Not Recorded	1249	1163 (57.8%)	82 (53.2%)	4 (25%)	.024*
<b>Pre-admission Elixhauser score, Mean (SD) <sup>3</sup></b>	2.2 (4.5)	0 (0)	4.4 (9.2)	2.5 (3.6)	.074
<b>Pre-admission Conditions (Selected), N (%) <sup>4</sup></b>					
Alcohol abuse	212	186 (9.2 %)	25 (15.3 %)	1 (6.2 %)	<.001*
Drug abuse	207	190 (9.4 %)	16 (9.8 %)	1 (6.2 %)	<.001*
Weight loss	123	109 (5.4 %)	13 (8 %)	1 (6.2 %)	<.001*
Cardiac arrhythmias	109	94 (4.7 %)	12 (7.4 %)	3 (18.8 %)	<.001*
<b>Number of pre-admission CNS codes, Mean (SD) <sup>3</sup></b>	0.2 (0.7)	0.1 (0.2)	1.9 (3.9)	0 (0)	.211
<b>Number of pre-admission PNS codes, Mean (SD) <sup>3</sup></b>	0 (0)	0 (0)	0 (0)	0.2 (0.6)	.067
<b>Severe Status, N (%)</b>					<.001*
Non-Severe	1674	1590 (78.5%)	72 (48.6%)	12 (70.6%)	
Severe	517	436 (21.5%)	76 (51.4%)	5 (29.4%)	
<b>Time to severe status, Mean days (SD) <sup>3</sup></b>	2.4 (5.3)	3.8 (9.5)	10.8 (33.1)	0.9 (2.5)	.269
<b>Survival, N (%)</b>					<.001*
Alive	2164	2001 (99%)	145 (94.2%)	18 (100%)	
Deceased	29	20 (1%)	9 (5.8%)	0 (0%)	
<b>Time to death, Mean days (SD) <sup>3</sup></b>	12.6 (36.3)	53.5 (76.3)	5.8 (10.2)	95 (268.7)	.269
<b>Time to discharge, Mean days (SD) <sup>3</sup></b>	5.1 (4.2)	3.8 (4.2)	5.8 (2.7)	5.9 (4.5)	.082
<b>Readmission, N (%)</b>					.957
Not Readmitted	1844	1707 (84.8%)	123 (83.7%)	14 (82.4%)	
Readmitted	332	305 (15.2%)	24 (16.3%)	3 (17.6%)	
<b>Time to first readmission, Mean days (SD) <sup>3</sup></b>	20 (20.2)	20.1 (19)	75.3 (86)	14.4 (22.8)	.171
<b>Number of readmissions, Mean (SD) <sup>3</sup></b>	0.1 (0.3)	0 (0)	0.2 (0.4)	0.4 (0.5)	.082

Notes:

1. Neurological status during acute COVID-19 hospitalization: NNC=No Neurological Condition; CNS=Central Nervous System diagnosis; PNS=Peripheral Nervous System diagnosis.
2. P-values were adjusted with the Benjamini-Hochberg method to control the false discovery rate when evaluating the distribution (categorical variables) or means (continuous variables) of characteristics stratified by neurological status. P-values < .05 were deemed significant and designated with asterisk (\*).
3. Continuous variables reflect the overall cohort mean and standard deviation (SD) of the median values reported by each healthcare system.
4. We report the top 4 most frequent pre-admission health conditions (*i.e.*, comorbidities) in the pediatric population. Please see S7 Table for all pre-admission health conditions stratified by neurological status during acute COVID-19 hospitalization.